

The effect of turbidity on the transition from social to sexual behavior in an African cichlid fish

Taylor K. Hrabak, Richard C. Oldham, and Dr. Suzanne M. Gray



INTRODUCTION

- Aquatic ecosystems are increasingly affected by human-induced stressors (e.g. elevated turbidity)^{1,2}.
- Light scattered by particles in the water column will alter the effectiveness of visual signals^{3,4}.
- *Pseudocrenilabrus multicolor victoriae* (a widespread African cichlid fish; Fig. 2) rely on visual cues such as male coloration and courtship displays for reproduction^{1,4}.
- To compensate for environmental stressors, fish must adapt behaviorally, morphologically, and physiologically, or face extinction^{1,2,4}.
- Evidence suggests that fish in turbid water often exhibit behavioral energetic trade-offs to persist^{2,5,6}.




Figure 1. Lwamunda Swamp (star) borders much of Lake Nabugabo, a satellite lake of Lake Victoria, Uganda, Africa.

OBJECTIVE

- To determine if exposure to elevated turbidity will alter the developmental transition from ecological behaviors (e.g. feeding, social) to reproductive behaviors (e.g. courtship displays) in the ontogenesis of *P. multicolor* (Fig. 2).




Figure 2. A juvenile *P. multicolor* (left) will begin displaying predominantly reproductive behaviors, rather than ecological behaviors, as an adult (right).

PREDICTION

- We predicted fish reared under turbid conditions would switch from ecological to reproductive behaviors earlier than those reared under clear conditions.

METHODS

- Ten independent broods of F1 *P. multicolor*, parental population originating from Lwamunda Swamp, Uganda, Africa (Fig. 1), were reared.
- Broods were split two weeks post-hatch and assigned to either a clear or turbid treatment (10 broods x 2 treatments= 20 tanks) (Fig. 3).
- Turbid tanks were maintained at ~10 NTU to mimic conditions in natural rivers.
- Beginning at three weeks post-hatch, each tank was filmed once a week for 20 min. (10 min. acclimation + 10 min. observation).
- Filming was concluded at 20 weeks of age (considered avg. age of sexual maturity in *P. multicolor*³).
- A subset of videos (n=75) were analyzed by performing focal follows on one male/tank and enumerating ecological and reproductive behaviors performed for the proportion of time active.




Figure 3. Half of each brood was reared under A) Clear conditions (~ 0 NTU) and the other half under B) Turbid conditions (~ 10 NTU).

RESULTS

- Fish in clear and turbid tanks spent the same amount of time being active (Fig. 4).
- With age, fish in both treatments performed fewer ecological behaviors and more reproductive behaviors; however, fish in turbid tanks performed more feeding behaviors overall (Fig. 5).
- The proportion of sexual behaviors, relative to total behaviors, increased with age in both treatments; however, fish in clear tanks performed significantly more sexual behaviors than those in turbid tanks (Fig. 6).
- Fish in clear tanks exhibited a shift from ecological behaviors to reproductive behaviors significantly earlier (8.5 weeks) than turbid-reared fish (12 weeks) (Fig. 7).




Figure 4. Mean (\pm SE) proportion of time spent active to inactive in fish reared under turbid and clear treatments ($F_{1,73} = 0.613$, $p = 0.436$).




Figure 5. Proportion of feeding behaviors to total behaviors in clear and turbid tanks over 18 weeks (Clear: Pearson's $r = -0.508$, $t_{38} = -3.638$, $p < 0.001$; Turbid: Pearson's $r = -0.398$, $t_{33} = -2.497$, $p = 0.017$).




Figure 6. Proportion of sexual behaviors to total behaviors in clear and turbid tanks over 18 weeks (Clear: Pearson's $r = 0.758$, $t_{38} = 7.175$, $p < 0.001$; Turbid: Pearson's $r = 0.688$, $t_{33} = 5.448$, $p < 0.001$).




Figure 7. The transition from ecological (<10%) and reproductive (>11%) over 18 weeks (Clear: $z_{1,39} = 3.42$, $p = 0.0062$; Turbid: $z_{1,34} = 2.941$, $p = 0.003$).

CONCLUSION

- Exposure to a novel environmental stressor, such as turbidity, may cause fish to reproduce later in their development.
- This may have various detrimental effects on the overall fitness and persistence of *P. multicolor* populations^{7,8}.
- Ultimately, this could pose a threat to the structure and function of Lake Victoria's food web^{7,8}.

LITERATURE CITED: 1. Gray et al. 2011. *Ecol Freshw Fish* 20: 259-236. 2. Reid et al. 2018. *Biol. Rev.* 3. Gray et al. 2012. *Curr Zool* 58:146-157. 4. Seehausen et al. 1997. *Science* 277:1808-1811. 5. Candolin, U. and J. Heuschele. 2008. *Trends Ecol Evol* 23:446-452. 6. Järvenpää, M. and K. Lindström. 2004. *Proc R Soc Lond [Biol]* 271:2361-2365. 7. Seehausen et al. 2008. *Nature* 455:620-626. 8. Henley et al. 2002. *Rev Fish Sci* 8:125-139.

ACKNOWLEDGEMENTS: Special thanks to Dr. Suzanne Gray, without whom this project would have not been possible. Additional thanks to Richard C. Oldham, Tiffany L. Atkinson, Chelsey L. Niemen, Brynne Drohan, and Nate Steffensmeier for their dedication to assisting me throughout each step of this process.